

HOLOMORPHIC EXTENSION FROM THE SPHERE TO THE BALL

LUCA BARACCO

ABSTRACT. Real analytic functions on the boundary of the sphere which have separate holomorphic extension along the complex lines through a boundary point have holomorphic extension to the ball. This was proved in [2] by an argument of CR geometry. We give here an elementary proof based on the expansion in holomorphic and antiholomorphic powers.

1. MAIN RESULT - STATEMENT AND PROOF

In my note [2] I stated a general principle of holomorphic extension from a convex boundary of $\mathbb{C}^n$ for functions which have separate holomorphic extension along generic $(2n - 2)$ -parameter families of discs. In particular, the discs which pass through a fixed boundary point. The conclusion seems to be natural in the framework of CR geometry but less in the classical literature of harmonic analysis. Also, it is in contrast with the case in which the “center” of the system of discs is an interior point of the ball: a single point does not suffice. For this reason, I want to supply here a direct proof which only uses Taylor expansions: moment condition forces coefficients of the antiholomorphic powers to vanish. The technique is specific of the sphere. General convex domains are out of reach unless one brings into play, as I do in [2], the extraordinary strength of the Riemann-Lempert Theorem (cf. [4] and [3]). This makes equivalent the family of discs by a point of a convex domain to the lines by a point of the ball.

Let $\mathbb{B}^n$ be the ball in $\mathbb{C}^n$, $\partial\mathbb{B}^n$ the sphere, z_o a point in $\partial\mathbb{B}^n$, and C^ω the real analytic functions.

Theorem 1.1. *Let f be a function in $C^\omega(\partial\mathbb{B}^n)$ and suppose that f extends from $\partial\mathbb{B}^n$ along each line passing through z_o . Then f extends holomorphically to $\mathbb{B}^n$.*

Proof. (a) We first prove the result for $\mathbb{B}^2$ in $\mathbb{C}^2$. It is not restrictive that z_o is the pole $(0, 1)$. The straight discs through $(0, 1)$ can be parametrized over $a \in \mathbb{C}$ as the sets D_a described by

$$D_a(\tau) = \left(\frac{\tau - 1}{1 + |a|^2} a, \frac{\tau - 1}{1 + |a|^2} + 1 \right) \quad \forall \tau \in \bar{\Delta}.$$

Note that when $|a| \gg 1$ the disc D_a becomes very close to the complex tangent line to the sphere at the point z_o , and moreover D_a lies in a neighborhood of z_o .

Since $f \in C^\omega(\partial\mathbb{B}^2)$, and $\bar{\partial}_{z_2}$ is transverse to $\partial\mathbb{B}^2$ at z_o , f can be extended in a neighborhood of z_o holomorphically in z_2 . We denote again by f this extension. We consider the power series development of f at z_o

$$f(z_1, \bar{z}_1, z_2) = \sum_{l=0}^{+\infty} \sum_{h+k+2m=l} b_{h,k,m} z_1^h \bar{z}_1^k z_2^m$$

note that we reordered the terms in a weighted degree (giving weight 2 to z_2). Taking $|a|$ sufficiently big we consider the N -momentum on the disc D_a :

$$(1.1) \quad G(a, N) = \int_{\partial\Delta} \tau^N f(D_a(\tau)) d\tau$$

$$(1.2) \quad = \int_{\partial\Delta} \tau^N \sum_{l=0}^{+\infty} \sum_{h+k+2m=l} b_{h,k,m} \left(\frac{\tau-1}{1+|a|^2} a \right)^h \left(\frac{\tau-1}{1+|a|^2} \right)^k \left(\frac{\tau-1}{1+|a|^2} \right)^m$$

We want to prove that $b_{h,k,m} = 0$ whenever $k > 0$. To this end, let l_o be the lowest weighted degree such that $b_{h,k,m} \neq 0$ for some $k > 0$ and let k_o be the highest degree in $\bar{z}_1$ for which this happens. We get $G(a, N) = 0$ for any N and any a , in particular, for ta with $|a| = 1$ and $t \rightarrow +\infty$. Consider the limit

$$(1.3)$$

$$(1.4) \quad \lim_{t \rightarrow +\infty} G(ta, N) t^{l_o} = \lim_{t \rightarrow +\infty} \sum_{l=l_o}^{+\infty} \sum_{h+k+2m=l, k>N} 2\pi i (-1)^{k+h+N-1} \times$$

$$\times \binom{h+k+m}{k-N-1} a^h \bar{a}^k \left(\frac{1}{t^2} + |a|^2 \right)^m t^{l_o-l} \frac{1}{\left(\frac{1}{t^2} + |a|^2 \right)^l}$$

$$(1.5) \quad = \sum_{h+k+2m=l_o, k>N} (2\pi i) (-1)^{h+k+N-1} \binom{h+k+m}{k-N-1} b_{h,k,m} \frac{a^h \bar{a}^k |a|^{2m}}{|a|^{2l_o}} = 0,$$

where we have used the fact that $\int_{\partial\Delta} \tau^N (\tau-1)^h (\bar{\tau}-1)^k (\tau-1)^m d\tau = \int_{\partial\Delta} (-1)^k \frac{\tau-1)^{h+k+m}}{\tau^{(k-l)}} d\tau = (-1)^{h+m+N+1} \binom{h+k+m}{k-1-N}$. Now, choosing $N = k_o - 1$, we get the following relation on the coefficients b 's:

$$\sum_{h+k_o+2m=l_o} (-1)^{h+m+k_o} \binom{h+k_o+m}{1} b_{h,k_o,m} a^{h+m} \bar{a}^{k_o+m} = 0.$$

Writing $a = e^{i\theta}$, we get

$$\sum_{h+k_o+2m=l_o} (-1)^{h+m+k_o} \binom{h+k_o+m}{1} b_{h,k_o,m} e^{i\theta(h-k_o)} = 0,$$

which implies $b_{h,k_o,m} = 0$ for $h+k_o+2m=l_o$. Therefore, when $k \geq 1$, we have $b_{h,k,m} = 0$ for any weighted degree l . This concludes the proof in dimension 2.

(b) We pass to the ball $\mathbb{B}^n$ of general dimension. We still suppose that z_o is the pole $(0, \dots, 1)$. By (a) we know that f extends holomorphically along the slices of $\mathbb{B}^n$ with the 2-dimensional planes through 0 and z_o . The different extensions glue to a single, well defined, function on $\mathbb{B}^n$ because, on the complex line L through 0 and z_o where they all overlap, they have to coincide by the Cauchy formula over $L \cap \partial\mathbb{B}^n$. Also, by Cauchy-Kowalevski Theorem, f extends to a neighborhood of z_o in $\mathbb{B}^n$ as a holomorphic function of z_n . This must coincide with the extension along the slices but it is, in addition, C^ω . In any of these slices f is holomorphic in radial directions and therefore, by Forelli's Theorem (cf. for instance [6]), it is holomorphic. Since it is also holomorphic in z_n , then it is holomorphic in a neighborhood of z_o in $\mathbb{B}^n$. By real analyticity, the holomorphicity of the extension propagates from a neighborhood of z_o to a neighborhood of the whole $\partial\mathbb{B}^n$. Finally, by Hartogs' Theorem, f is in fact holomorphic in the whole $\mathbb{B}^n$. □

REFERENCES

- [1] **M. Agranovsky**—Holomorphic extension from the unit sphere in $\mathbb{C}^n$ into complex lines passing through a finite set *math. arXiv 0910.3592* (2009)
- [2] **L. Baracco**—Holomorphic extension from a convex hypersurface, (2009)
- [3] **C.H. Chang, M.C. Hu and H.P. Lee**—Extremal analytic discs with prescribed boundary data, *Trans. of the A.M.S.*, **310** 1 (1988), 355–369
- [4] **L. Lempert**—La métrique de Kobayashi et la représentation des domaines sur la boule. (French) [The Kobayashi metric and the representation of domains on the ball *Bull. Soc. Math. France* **109** (1981), 427-474
- [5] **L. Lempert**—A precise result on the boundary regularity of biholomorphic mappings, *Math. Z.* **193** (1986), 559–579
- [6] **W. Rudin**—Function theory on the unit ball of $\mathbb{C}^n$, Grundlehren Math. W., Springer Verlag New York - Berlin (1980)

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI PADOVA, VIA TRIESTE 63, 35121 PADOVA, ITALY
E-mail address: baracco@math.unipd.it